

Getting started with XOSPATA™ (gilteritinib)

	Pre-treatment	Month	1			2	3	4	5	6	>6 months
		Day	1	8	15						
Testing and Assessments											
FLT3-ITD/FLT3-TKD MUTATION STATUS¹ (before taking XOSPATA, all patients must have confirmation of FLT3-ITD/FLT3-TKD mutation using a validated test)											
BLOOD CHEMISTRIES, INCLUDING CPK¹ (before initiation of treatment, on Day 15, and monthly for the duration of treatment. If present, correct hypokalaemia or hypomagnesaemia prior to and during treatment)											
ECG¹ (before initiation of treatment, on Days 8 and 15 of Cycle 1 and prior to the start of the next three subsequent months of treatment)											
PREGNANCY¹ Females of reproductive potential should be advised to have a pregnancy test within seven days prior to starting treatment with XOSPATA and to use effective contraception during treatment with XOSPATA and for at least 6 months after stopping treatment.											
Dosing											
STARTING DOSE & DURATION¹ The recommended starting dose is 120 mg (three 40 mg tablets) once daily	3 x 40 mg OD Continue treatment until patient is no longer clinically benefiting or until unacceptable toxicity occurs. Consider treatment at the prescribed dose for up to 6 months to allow time for a clinical response										
DOSE ESCALATION IF REQUIRED¹	5 x 40 mg OD If patient does not achieve a CRc after 4 weeks consider increasing dose to 200 mg (5 x 40 mg) once daily if tolerated or clinically warranted. Consider treatment for up to 6 months to allow time for a clinical response										
Adverse Event Management											
POSTERIOR REVERSIBLE ENCEPHALOPATHY SYNDROME	Discontinue XOSPATA										
QT PROLONGATION ON DAY 8 OF CYCLE 1¹ QTc interval increased by >30 msec on ECG on Day 8 of Cycle 1	Confirm with ECG on Day 9. If confirmed, consider dose reduction to 80 mg [†]										
QTcF INTERVAL >500 MSEC	Interrupt XOSPATA. Resume XOSPATA at a reduced dose (80 mg or 120 mg [†]) when QTcF interval returns to within 30 msec of baseline or ≤480 msec										
DIFFERENTIATION SYNDROME¹	If differentiation syndrome is suspected, administer corticosteroids and initiate hemodynamic monitoring. Interrupt XOSPATA if severe signs and/or symptoms persist for more than 48 hours after initiation of corticosteroids. Resume XOSPATA at the same dose when signs and symptoms improve to Grade 2 [‡] or lower.										
PANCREATITIS¹	Interrupt XOSPATA until pancreatitis is resolved. Resume treatment with XOSPATA at a reduced dose (80 mg or 120 mg [†]).										
OTHER GRADE 3[‡] OR HIGHER AEs RELATED TO XOSPATA¹	Interrupt XOSPATA until toxicity resolves or improves to Grade 1 [‡] . Resume treatment with XOSPATA at a reduced dose (80 mg or 120 mg [†]).										

Please consult the Summary of Product Characteristics for detailed information on testing, posology and management of adverse events.

XOSPATA is indicated as monotherapy for the treatment of adult patients who have relapsed or refractory acute myeloid leukaemia (AML) with a FLT3 mutation.

* The daily dose can be reduced from 120 mg to 80 mg or from 200 mg to 120 mg.¹

† If XOSPATA is re-introduced at a reduced dose, ECG should be performed after 15 days of dosing and prior to start of the next three subsequent months of treatment.¹

‡ Grade 1 is mild, Grade 2 is moderate, Grade 3 is serious, Grade 4 is life-threatening.¹

XOSPATA™
gilteritinib 40mg tablets

ABBREVIATED PRESCRIBING INFORMATION OF XOSPATA™ 40 mg film-coated tablets

(Please refer the full prescribing information for further details) **1. Name of the medicinal product:** XOSPATA™ 40 mg film-coated tablets **2. QUALITATIVE AND QUANTITATIVE COMPOSITION:** Each film-coated tablet contains 40 mg gilteritinib (as fumarate). **3. PHARMACEUTICAL FORM:** Film-coated tablet (tablet). **4. CLINICAL PARTICULARS** **4.1 Therapeutic indication:** Xospata is indicated as monotherapy for the treatment of adult patients who have relapsed or refractory acute myeloid leukaemia (AML) with a FLT3 mutation. **4.2 Posology and method of administration** Posology: The recommended starting dose is 120 mg gilteritinib (three 40 mg tablets) once daily. Blood chemistries and ECG monitoring should be performed before initiation and during the treatment as recommended. Treatment should continue until the patient is no longer clinically benefiting from Xospata or until unacceptable toxicity occurs. Response may be delayed; therefore, continuation of treatment at the prescribed dose for up to 6 months should be considered to allow time for a clinical response. In the absence of a response after 4 weeks of treatment, the dose can be increased to 200 mg once daily, if tolerated or clinically warranted. **Dose modifications:** Xospata should be administered at about the same time each day. If a dose is missed or not taken at the usual time, the dose should be administered as soon as possible on the same day, and patients should return to the normal schedule the following day. If vomiting occurs after dosing, patients should not take another dose but should return to the normal schedule the following day. **Elderly:** No dose adjustment is required in patients ≥65 years of age. **Hepatic impairment:** No dose adjustment is required for patients with mild or moderate hepatic impairment. Xospata is not recommended for use in patients with severe hepatic impairment, as safety and efficacy have not been evaluated in this population. **Renal impairment:** No dose adjustment is necessary in patients with mild or moderate renal impairment. There is no clinical experience in patients with severe renal impairment. **Paediatric population:** The safety and efficacy of Xospata in children aged below 18 years has not yet been established. No data is available. Due to in vitro binding to 5HT2B, there is a potential impact on cardiac development in patients less than 6 months of age. **4.3 Contraindications:** Hypersensitivity to the active substance or to any of the excipients listed in section 6.1 (Kindly refer full prescribing information). **4.4 Special warnings and precautions for use:** **Differentiation syndrome:** If differentiation syndrome is suspected, corticosteroid therapy should be initiated along with hemodynamic monitoring until symptom resolution. If severe signs and/or symptoms persist for more than 48 hours after initiation of corticosteroids, Xospata should be interrupted until signs and symptoms are no longer severe. Corticosteroids can be tapered after resolution of symptoms and should be administered for a minimum of 3 days. Symptoms of differentiation syndrome may recur with premature discontinuation of corticosteroid treatment. **Posterior reversible encephalopathy syndrome:** There have been reports of posterior reversible encephalopathy syndrome (PRES) in patients receiving Xospata (see section 4.6). If PRES is suspected, it should be confirmed by brain imaging, preferably magnetic resonance imaging (MRI). Discontinuation of Xospata in patients who develop PRES is recommended. **Prolonged QT interval:** QT prolongation can be observed in the first three months of treatment with gilteritinib. Therefore, electrocardiogram (ECG) should be performed prior to initiation of treatment, on day 8 and 15 of cycle 1, and prior to the start of the next three subsequent months of treatment. Caution is warranted in patients with relevant cardiac history. Hypokalaemia or hypomagnesaemia may increase the QT prolongation risk. Hypokalaemia or hypomagnesaemia should therefore be corrected prior to and during Xospata treatment. Xospata should be interrupted in patients who have a QTcF >500 msec. The decision to re-introduce gilteritinib treatment after an event of QT prolongation should be based on a careful consideration of benefits and risks. If Xospata is re-introduced at a reduced dose, ECG should be performed after 15 days of dosing, and prior to the start of the next three subsequent months of treatment. In clinical studies, 12 patients had QTcF >500 msec. Three patients interrupted and re-initiated treatment without recurrence of QT prolongation. **Pancreatitis:** Xospata should be interrupted and can be resumed at a reduced dose when the signs and symptoms of pancreatitis have resolved. **Interactions:** Co-administration of CYP3A/P-gp inducers may lead to decreased gilteritinib exposure and consequently a risk for lack of efficacy. Therefore, concomitant use of gilteritinib with strong CYP3A4/P-gp inducers should be avoided. Caution is required when concomitantly prescribing gilteritinib with medicinal products that are strong inhibitors of CYP3A, P-gp and/or breast cancer resistant protein (BCRP) because they can increase gilteritinib exposure. In situations where satisfactory therapeutic alternatives do not exist, patients should be closely monitored for toxicities during administration of gilteritinib. Concomitant use of gilteritinib with medicinal products that target 5HT2B receptor or sigma nonspecific receptors should be avoided unless use is considered essential for the care of the patient since gilteritinib may reduce the effect of these drugs. **Embryofoetal toxicity and contraception:** Pregnant women should be informed of the potential risk to a foetus. Females of reproductive potential should be advised to have a pregnancy test within seven days prior to starting treatment with Xospata and to use effective contraception during treatment with Xospata and for at least 6 months after stopping treatment. Women using hormonal contraceptives should add a barrier method of contraception. Males with female partners of reproductive potential should be advised to use effective contraception during treatment and for at least 4 months after the last dose of Xospata. **4.5 Effects on ability to drive and use machines:** Dizziness has been reported in patients taking Xospata and should be considered when assessing a patient's ability to drive or use machines. **4.6 Undesirable effects:** **Summary of the safety profile:** The safety of Xospata was evaluated in 319 patients with relapsed or refractory AML who have received at least one dose of 120 mg gilteritinib. The most frequent adverse reactions with gilteritinib were alanine aminotransferase (ALT) increased (82.1%), aspartate aminotransferase (AST) increased (80.6%), blood alkaline phosphatase increased (68.7%), blood creatine phosphokinase increased (53.9%), diarrhoea (35.1%), fatigue (30.4%), nausea (29.8%), constipation (28.2%), cough (28.2%), peripheral oedema (24.1%), dyspnoea (24.1%), dizziness (20.4%), hypotension (17.2%), pain in extremity (14.7%), asthenia (13.8%), arthralgia (12.5%) and myalgia (12.5%). The most frequent serious adverse reactions were acute kidney injury (6.6%), diarrhoea (4.7%), ALT increased (4.1%), dyspnoea (3.4%), AST increased (3.1%) and hypotension (2.8%). Other clinically significant serious adverse reactions included differentiation syndrome (2.2%), electrocardiogram QT prolonged (0.9%) and posterior reversible encephalopathy syndrome (0.6%). **OVERDOSE:** There is no known specific antidote for Xospata. In the event of an overdose, treatment with Xospata should be stopped. Patients must be closely monitored for signs or symptoms of adverse reactions, and appropriate symptomatic and supportive treatment initiated, taking into consideration the long half-life estimated at 113 hours. **For full prescribing information please write to:** **MARKETING AUTHORISATION HOLDER: Astellas Pharma India Private Limited**, 301, 3rd Floor, C and B Square, 127 Andheri Kurla Road, Chakala, Andheri (East), Mumbai - 400 069. (Version: XOSPATA/aPI/India/Ver.2.0/Apr2024).

Reference: 1. XOSPATA Summary of Product Characteristics.

AE – adverse event; CPK – creatine phosphokinase; CRc – composite complete remission; ECG – electrocardiogram; FLT3-ITD – FMS-like tyrosine kinase 3 internal tandem duplication; FLT3-TKD – FMS-like tyrosine kinase 3 tyrosine kinase domain; OD – once-daily; PRES – posterior reversible encephalopathy syndrome; QTc – QT corrected

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only, Additional Information Available on request.



Astellas Pharma India Private Limited

Unit No. 301, C & B Square, CTS Number 95 A, 127 Andheri Kurla Road, Chakala, Andheri East, Mumbai, Maharashtra 400 069.

Tel: 022 61557600 | www.astellas.com | Email: pv@in.astellas.com

Date of preparation: September 2025 MAT-IN-XOS-2025-00064

Scan for ABPI



XOSPATA™
gilteritinib 40mg tablets